# LECTURE 3: FANO'S INEQUALITY AND APPLICATIONS

### 1 Introduction

So far, we have focused on problems concerning the *sufficiency* of sample sizes. For instance, we might ask for the minimum number of samples n required for an algorithm like Empirical Risk Minimization (ERM) to work with a certain guarantee.

Now, we shift our perspective to ask a different question: what happens if n is not sufficient? We want to determine the *necessary* number of samples for *any* learning algorithm to succeed. This involves establishing lower bounds on the sample complexity.

The general setting we consider is as follows:

- 1. Nature picks a "true" hypothesis  $\tilde{f}$  from a finite hypothesis class  $\mathcal{H}$ .
- 2. A dataset S of size n is generated, conditioned on the choice of  $\tilde{f}$ .
- 3. A learner observes the dataset S and produces an estimate  $\hat{f} \in \mathcal{H}$ . This process defines a Markov Chain:  $\tilde{f} \to S \to \hat{f}$ .

Our goal is to understand the conditions under which the probability of making a mistake,  $\mathbb{P}\left[\hat{f} \neq \tilde{f}\right]$ , is high, regardless of the learning algorithm used.

# 2 Information Theory Basics

#### 2.1 Entropy

**Definition 1** (Entropy). The entropy of a discrete random variable X with support  $\mathcal{X}$  and probability mass function p(x) is defined as:

$$H(X) = -\sum_{x \in \mathcal{X}} p(x) \log(p(x)).$$

Entropy measures the average uncertainty of a random variable. It has the following properties:

- 1.  $H(X) \ge 0$ .
- 2.  $H(X) \leq \log |\mathcal{X}|$ .

*Proof of property 2.* We use Jensen's inequality, which states that for a concave function  $\phi$ , we have

 $\mathbb{E}[\phi(Y)] \leq \phi(\mathbb{E}[Y])$ . The logarithm function is concave.

$$H(X) = -\sum_{x \in \mathcal{X}} p(x) \log(p(x)) = \sum_{x \in \mathcal{X}} p(x) \log\left(\frac{1}{p(x)}\right) = \mathbb{E}_{X \sim p} \left[\log\left(\frac{1}{p(X)}\right)\right]$$

$$\leq \log\left(\mathbb{E}_{X \sim p} \left[\frac{1}{p(X)}\right]\right) \quad \text{(by Jensen's inequality)}$$

$$= \log\left(\sum_{x \in \mathcal{X}} p(x) \frac{1}{p(x)}\right) = \log\left(\sum_{x \in \mathcal{X}} 1\right) = \log|\mathcal{X}|.$$

**Definition 2** (Conditional Entropy). *The conditional entropy of a random variable* Y *given* X *is defined as:* 

$$H(Y|X) = \sum_{x \in \mathcal{X}} \mathcal{P}_X(x) H(Y|X = x)$$

$$= -\sum_{x \in \mathcal{X}} \mathcal{P}_X(x) \sum_{y \in \mathcal{Y}} \mathcal{P}_{Y|X}(y|x) \log \mathcal{P}_{Y|X}(y|x)$$

$$= -\sum_{x \in \mathcal{X}, y \in \mathcal{Y}} \mathcal{P}_{XY}(x, y) \log \mathcal{P}_{Y|X}(y|x).$$

Fact 1 (Chain Rule for Entropy). The joint entropy of two random variables X and Y can be expressed as:

$$H(X,Y) = H(X) + H(Y|X).$$

Similarly, for three variables: H(X,Y|Z) = H(X|Z) + H(Y|X,Z).

#### 2.2 Mutual Information

**Definition 3** (Mutual Information). *The mutual information between two random variables* X *and* Y *is defined as:* 

$$I(X,Y) = \sum_{x \in \mathcal{X}, y \in \mathcal{Y}} \mathcal{P}_{XY}(x,y) \log \frac{\mathcal{P}_{XY}(x,y)}{\mathcal{P}_{X}(x)\mathcal{P}_{Y}(y)}.$$

Mutual information measures the reduction in uncertainty about one random variable given knowledge of another. It has the following key properties:

- 1. I(X,Y) > 0.
- 2. I(X,Y) = 0 if and only if X and Y are independent.
- 3. I(X,Y) = H(X) H(X|Y).

**Fact 2** (Conditioning Reduces Entropy). For any two random variables X and Y, we have:

$$H(X|Y) < H(X)$$
.

This follows directly from the properties I(X,Y) = H(X) - H(X|Y) and  $I(X,Y) \ge 0$ .

Instructor: Adarsh Barik
Scribed by: Akshat Jha

Lecture 3, COV878
August 21, 2025

**Definition 4** (Conditional Mutual Information). *The mutual information between X and Y conditioned on a third variable Z is:* 

$$I(X,Y|Z) = H(X|Z) - H(X|Y,Z).$$

Fact 3 (Chain Rule for Mutual Information).

$$I(X,(Y,Z)) = I(X,Y) + I(X,Z|Y).$$

*Proof of fact 3.* The definition of mutual information is I(X,Y) = H(X) - H(X|Y). The definition of conditional mutual information is I(X,Y|Z) = H(X|Z) - H(X|Y,Z).

RHS = 
$$I(X,Y) + I(X,Z|Y)$$
  
=  $[H(X) - H(X|Y)] + [H(X|Y) - H(X|Y,Z)]$  (substituting the definitions)  
=  $H(X) - H(X|Y) + H(X|Y) - H(X|Y,Z)$   
=  $H(X) - H(X|Y,Z)$  (canceling terms)  
=  $I(X,(Y,Z)) = LHS$  (by the definition of mutual information)

# 3 Markov Chains and the Data Processing Inequality

**Definition 5** (Markov Chain). *Random variables X,Y,Z are said to form a Markov Chain, denoted X*  $\rightarrow$  *Y*  $\rightarrow$  *Z, if their joint probability distribution can be written as:* 

$$\mathcal{P}_{XYZ}(x,y,z) = \mathcal{P}_X(x)\mathcal{P}_{Y|X}(y|x)\mathcal{P}_{Z|Y}(z|y)$$
.

This is equivalent to the statement that X and Z are conditionally independent given Y, which implies I(X, Z|Y) = 0.

**Theorem 1** (Data Processing Inequality). *If*  $X \to Y \to Z$  *form a Markov Chain, then* 

$$I(X,Z) \leq I(X,Y)$$
.

Intuitively, no amount of processing on Y (to get Z) can increase the information that Y contains about X.

*Proof.* By the chain rule for mutual information, we have two ways to expand I(X, (Y, Z)):

$$I(X,(Y,Z)) = I(X,Y) + I(X,Z|Y)$$
  
=  $I(X,Z) + I(X,Y|Z)$ 

Since  $X \to Y \to Z$  is a Markov chain, we know I(X, Z|Y) = 0. Therefore:

$$I(X,Y) = I(X,Z) + I(X,Y|Z)$$

Because mutual information is non-negative,  $I(X,Y|Z) \ge 0$ . Thus, we conclude that  $I(X,Y) \ge I(X,Z)$ .

 $\stackrel{}{\neg}$ 

Instructor: Adarsh BarikLecture 3, COV878Scribed by: Akshat JhaAugust 21, 2025

# 4 Fano's Inequality

Fano's inequality provides a lower bound on the probability of error for any estimator in a classification problem. It connects the probability of error with the conditional entropy of the true hypothesis given the data.

**Theorem 2** (Fano's Inequality). Consider the learning setup where Nature picks a true hypothesis  $\tilde{f} \in \mathcal{H}$ , data S is generated conditioned on  $\tilde{f}$ , and a learner produces an estimate  $\hat{f}$  from S. For any such estimator  $\hat{f}$ , the probability of error  $\mathcal{P}_e = \mathbb{P}\left[\hat{f} \neq \tilde{f}\right]$  is bounded. A simplified and useful version of the inequality is:

$$\mathcal{P}_e \log |\mathcal{H}| \ge H(\tilde{f}|S) - \log(2)$$
.

**Corollary 1** (Simplified Fano's Inequality). Let the true hypothesis  $\tilde{f}$  be chosen uniformly at random from the hypothesis class  $\mathcal{H}$ . Then  $H(\tilde{f}) = \log |\mathcal{H}|$ . Using the relation  $H(\tilde{f}|S) = H(\tilde{f}) - I(\tilde{f},S)$ , we can rearrange the inequality to get:

$$\mathbb{P}\left[\hat{f} \neq \tilde{f}\right] \geq 1 - \frac{I(\tilde{f}, S) + \log 2}{\log |\mathcal{H}|}.$$

This form is particularly useful. To get a high probability of error (i.e., a lower bound close to 1), we need to show that the mutual information  $I(\tilde{f}, S)$  is small compared to  $\log |\mathcal{H}|$ .

# 5 A Lower Bound on Sample Complexity

Our goal is to construct a learning problem where any algorithm must fail. To do this using Fano's inequality, we need to find an upper bound on the mutual information  $I(\tilde{f}, S)$ .

### 5.1 Kullback-Leibler (KL) Divergence

**Definition 6** (KL Divergence). Let  $\mathcal{P}$  and  $\mathcal{Q}$  be two probability distributions on the same support  $\mathcal{X}$ , with probability mass functions p(x) and q(x) respectively. The KL-divergence between  $\mathcal{P}$  and  $\mathcal{Q}$  is defined as:

$$KL(\mathcal{P}||\mathcal{Q}) = \sum_{x \in \mathcal{X}} p(x) \log \frac{p(x)}{q(x)}.$$

**Fact 4** (Additivity of KL Divergence). Let  $\mathcal{P}_{XY}$  and  $\mathcal{Q}_{XY}$  be product distributions, i.e.,  $\mathcal{P}_{XY} = \mathcal{P}_X \mathcal{P}_Y$  and  $\mathcal{Q}_{XY} = \mathcal{Q}_X \mathcal{Q}_Y$ . Then:

$$\textit{KL}(\mathcal{P}_{XY}||\mathcal{Q}_{XY}) = \textit{KL}(\mathcal{P}_{X}||\mathcal{Q}_{X}) + \textit{KL}(\mathcal{P}_{Y}||\mathcal{Q}_{Y}) \; .$$

Instructor: Adarsh BarikLecture 3, COV878Scribed by: Akshat JhaAugust 21, 2025

### 5.2 Bounding Mutual Information via KL Divergence

The mutual information can be expressed in terms of KL divergence. When  $\tilde{f}$  is uniform over  $\mathcal{H}$ :

$$\begin{split} I(\tilde{f},S) &= \sum_{\tilde{f} \in \mathcal{H}} \sum_{S} \mathcal{P}_{\tilde{f},S}(\tilde{f},S) \log \frac{\mathcal{P}_{\tilde{f},S}(\tilde{f},S)}{\mathcal{P}_{\tilde{f}}(\tilde{f}) \, \mathcal{P}_{S}(S)} \\ &= \frac{1}{|\mathcal{H}|} \sum_{f \in \mathcal{H}} \sum_{S} \mathcal{P}_{S|f}(S) \log \frac{\mathcal{P}_{S|f}(S)}{\mathcal{P}_{S}(S)} \\ &= \frac{1}{|\mathcal{H}|} \sum_{f \in \mathcal{H}} \text{KL} \big( \mathcal{P}_{S|f} \, \| \, \mathcal{P}_{S} \big). \end{split}$$

Using the convexity of KL-divergence and expanding the  $\mathcal{P}_S$  in the denominator, one can further show that:

$$I(\tilde{f}, S) \leq \frac{1}{|\mathcal{H}|^2} \sum_{f \in \mathcal{H}} \sum_{f' \in \mathcal{H}} \mathrm{KL}(\mathcal{P}_{S|f} \parallel \mathcal{P}_{S|f'}).$$

Since the dataset S consists of n i.i.d. samples, by the additivity of KL divergence, this becomes:

$$I(\tilde{f}, S) \leq \frac{n}{|\mathcal{H}|^2} \sum_{f \in \mathcal{H}} \sum_{f' \in \mathcal{H}} KL(\mathcal{P}_{X,Y|f} \parallel \mathcal{P}_{X,Y|f'}).$$

#### 5.3 The Main Result

**Theorem 3.** Nature picks a "true" hypothesis  $\tilde{f}$  uniformly at random from  $\mathcal{H}$ . Then a dataset of n iid samples is generated conditioned on the choice of  $\tilde{f}$ . The learner infers  $\hat{f}$  from the data. There exists a specific learning problem and data distribution such that if the number of samples n satisfies:

$$n < \frac{\log(|\mathcal{H}|)/2 - \log 2}{48\epsilon^2}$$

for a fixed  $\epsilon \in (0, 1/8)$ , the learning fails, i.e.  $\mathbb{P}\left[\hat{f} \neq \tilde{f}\right] \geq \frac{1}{2}$ , for any mechanism that a learner could use for picking  $\hat{f}$ .

*Proof.* We construct a "hard" learning problem.

- 1. Constructing the hypothesis class: Let the input space be  $\mathcal{Z} = \{z_1, \ldots, z_d\}$ . For each vector  $\tau \in \{-1, 1\}^d$ , define a hypothesis  $h_\tau : \mathcal{Z} \to \{-1, 1\}$  such that  $h_\tau(z_i) = \tau_i$ . The hypothesis class is  $\mathcal{H} = \{h_\tau : \tau \in \{-1, 1\}^d\}$ , so  $|\mathcal{H}| = 2^d$ .
- 2. **Defining the data distribution**: Nature picks a  $\tau$  uniformly at random, setting the true hypothesis  $\tilde{f} = h_{\tau}$ . For each sample  $(x_i, y_i)$  in the dataset S:
  - $x_i$  is chosen uniformly from  $\mathcal{Z}$ .
  - The label  $y_i$  is a noisy version of the true label  $\tilde{f}(x_i)$ . Specifically, if  $x_i = z_i$ ,

$$\mathbb{P}\left[y_i = y | x_i = z_j, \tilde{f} = h_\tau\right] = \begin{cases} \frac{1}{2} + 2\epsilon & \text{if } y = \tau_j \\ \frac{1}{2} - 2\epsilon & \text{if } y = -\tau_j \end{cases}.$$

Instructor: Adarsh Barik
Scribed by: Akshat Jha

Lecture 3, COV878
August 21, 2025

3. **Bounding KL divergence**: For any two distinct hypotheses  $f, f' \in \mathcal{H}$ , we compute the KL divergence between their corresponding data distributions. Since  $P_{X|f}$  is uniform and independent of f, the KL divergence simplifies:

$$KL(\mathcal{P}_{X,Y|f} || \mathcal{P}_{X,Y|\bar{f}}) = \sum_{y \in \{-1,1\}} \sum_{x \in \mathcal{Z}} P_{x,y|f} \log \frac{P_{x,y|f}}{P_{x,y|\bar{f}}}$$

$$= \sum_{y \in \{-1,1\}} \sum_{x \in \mathcal{Z}} P_{x|f} P_{y|x,f} \log \frac{P_{y|x,f}}{P_{y|x,\bar{f}}}$$

$$= \frac{1}{d} \sum_{y \in \{-1,1\}} \sum_{x \in \mathcal{Z}} P_{y|x,f} \log \frac{P_{y|x,f}}{P_{y|x,\bar{f}}}$$

$$\leq \frac{d}{d} \sum_{y \in \{-1,1\}} P_{y|x=\tilde{x},f} \log \frac{P_{y|x=\tilde{x},f}}{P_{y|x=\tilde{x},\bar{f}}}$$

The last inequality is due to the fact that  $P_{y|x=\tilde{x},f}(\cdot)$  and  $P_{y|x=\tilde{x},\bar{f}}(\cdot)$  differ only when  $f(\tilde{x}) \neq \bar{f}(\tilde{x})$  for some  $\tilde{x} \in \mathcal{Z}$ . Also, notice that this can only happen for a maximum of d samples in  $\mathcal{Z}$ . Furthermore, due to our construction:

$$\sum_{y \in \{-1,1\}} P_{y|x=\tilde{x},f}(\cdot) \log \frac{P_{y|x=\tilde{x},f}(\cdot)}{P_{y|x=\tilde{x},\tilde{f}}(\cdot)} = \left(\frac{1}{2} + 2\epsilon\right) \log \left(\frac{\frac{1}{2} + 2\epsilon}{\frac{1}{2} - 2\epsilon}\right) + \left(\frac{1}{2} - 2\epsilon\right) \log \left(\frac{\frac{1}{2} - 2\epsilon}{\frac{1}{2} + 2\epsilon}\right)$$

$$\leq 48\epsilon^{2} \quad \text{for } \epsilon \in (0,1/8) \ .$$

4. **Bound mutual information**: Plugging this into our bound for  $I(\tilde{f}, S)$ :

$$I(\tilde{f}, S) \leq \frac{1}{|\mathcal{H}|^2} \sum_{f \in \mathcal{H}} \sum_{\tilde{f} \in \mathcal{H}} KL(\mathcal{P}_{S|f} || \mathcal{P}_{S|\bar{f}})$$

$$= \frac{n}{|\mathcal{H}|^2} \sum_{f \in \mathcal{H}} \sum_{\tilde{f} \in \mathcal{H}} KL(\mathcal{P}_{X,Y|f} || \mathcal{P}_{X,Y|\bar{f}}) \leq \frac{n}{|\mathcal{H}|^2} \sum_{f, f' \in \mathcal{H}} 48\epsilon^2 = n \cdot 48\epsilon^2.$$

5. Apply Fano's inequality:

$$\mathbb{P}\left[\hat{f} \neq \tilde{f}\right] \ge 1 - \frac{I(\tilde{f}, S) + \log 2}{\log |\mathcal{H}|}$$
$$\ge 1 - \frac{48n\epsilon^2 + \log 2}{d \log 2}.$$

We want this probability to be at least  $\frac{1}{2}$ .

$$1 - \frac{48n\epsilon^2 + \log 2}{d\log 2} > \frac{1}{2} \implies \frac{d\log 2}{2} > 48n\epsilon^2 + \log 2 \implies n < \frac{d\log(2)/2 - \log 2}{48\epsilon^2}$$

Since  $\log |\mathcal{H}| = d \log 2$ , this proves the theorem.

This result provides a fundamental limit on learnability, showing that if the sample size is too small relative to the complexity of the hypothesis class (measured by  $\log |\mathcal{H}|$ ) and the difficulty of the problem (inversely related to  $\epsilon$ ), no algorithm can be guaranteed to succeed.

Instructor: Adarsh BarikLecture 3, COV878Scribed by: Akshat JhaAugust 21, 2025

### 5.4 Connection to PAC Learning

The previous subsection demonstrated that it is possible to construct instances where the probability that  $\hat{f} \neq \tilde{f}$  remains bounded away from zero. However, this fact alone does not immediately imply a poor generalization (or risk) bound in the PAC framework. In this subsection, we provide an informal proof-sketch to connect the earlier result to the PAC setting.

We assume that nature picks the  $\tilde{f} = h_{\tau}$  as the "true" hypothesis. We denote the corresponding induced joint distribution on (x, y) as  $\mathcal{P}_{X,Y}^{\tau}$ . One can compute the risk of any hypothesis  $h \in \mathcal{H}$  as:

$$\begin{split} R(h) &= \mathbb{E}_{(x,y) \sim \mathcal{P}_{X,Y}^{\tau}} \left[ \mathbb{1}(h(x) \neq y) \right] \\ &= \sum_{y \in \{-1,1\}} \sum_{x \in \mathcal{Z}} \mathcal{P}_{X,Y}^{\tau}(x,y) \mathbb{1}(h(x) \neq y) \\ &= \sum_{y \in \{-1,1\}} \sum_{i=1}^{d} \mathcal{P}_{X}^{\tau}(z_{i}) \mathcal{P}_{Y|X=z_{i}}^{\tau}(y) \mathbb{1}(h(z_{i}) \neq y) \\ &= \frac{1}{d} \sum_{i=1}^{d} \left( \mathcal{P}_{Y|X=z_{i}}^{\tau}(y = \tau_{i}) \mathbb{1}(h(z_{i}) \neq \tau_{i}) + \mathcal{P}_{Y|X=z_{i}}^{\tau}(y = -\tau_{i}) \mathbb{1}(h(z_{i}) = \tau_{i}) \right) \\ &= (\frac{1}{2} - 2\epsilon) + \frac{4\epsilon}{d} \sum_{i=1}^{d} \mathbb{1}(h(z_{i}) \neq \tau_{i}) \end{split}$$

Notice that R(h) is minimized by picking  $h = h_{\tau}$ . Furthermore, one can also show that  $\sum_{i=1}^{d} \mathbb{1}(h(z_i) \neq \tau_i) = \Omega(d)$  in expectation.

Therefore, for any  $h \neq h_{\tau}$ ,

$$R(h) > R(h_{\tau}) + \Omega(\epsilon)$$
.

**Disclaimer:** These notes have not been scrutinized with the level of rigor usually applied to formal publications. Readers should verify the results before use.